



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 1990

Intravenous 1,25(OH)₂ Vitamin D₃ Therapy in Haemodialysis Patients: Evaluation of Direct and Calcium-Mediated Short-Term Effects on Serum Parathyroid Hormone Concentration

Probst, W ; Fischer, J A ; Binswanger, U

Abstract: Eleven patients on chronic haemodialysis treatment thrice weekly received 1 g 1,25(OH)₂D₃ i.v. after each dialysis for 3 weeks. Phosphate binders were mainly CaCO₃, supplemented in a few patients by moderate amounts of Al(OH)₃. Ionised calcium was measured by ion-selective electrode, normal values being 1.28-1.42mmol/l. PTH was estimated by an N-terminal-sensitive assay; normal values are <0.25 g/ml. Results before and after 1,25(OH)₂D₃ were: ionised calcium before haemodialysis, 1.19±0.12 and 1.17±0.14; ionised calcium after haemodialysis, 1.33±0.07 and 1.30±0.09; PTH before haemodialysis, 1.39±0.71 and 1.38±0.69; PTH after haemodialysis, 0.64±0.22 and 0.60±0.17; Phosphate before haemodialysis, 1.85±0.48 and 2.18±0.43 (P<0.05). No change of PTH concentration and ionised calcium before and after haemodialysis treatment could be documented after i.v. 1,25(OH)₂D₃ treatment. Mild and severe hyperparathyroidism were indistinguishable. Increased serum calcium concentrations therefore appear to be required for the suppression of PTH secretion by i.v. 1,25(OH)₂D₃ therapy

DOI: <https://doi.org/10.1093/ndt/5.6.457>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-154843>

Journal Article

Published Version

Originally published at:

Probst, W; Fischer, J A; Binswanger, U (1990). Intravenous 1,25(OH)₂ Vitamin D₃ Therapy in Haemodialysis Patients: Evaluation of Direct and Calcium-Mediated Short-Term Effects on Serum Parathyroid Hormone Concentration. *Nephrology, Dialysis, Transplantation*, 5(6):457-460.

DOI: <https://doi.org/10.1093/ndt/5.6.457>

Original Article

Intravenous 1,25(OH)₂ Vitamin D3 Therapy in Haemodialysis Patients: Evaluation of Direct and Calcium-Mediated Short-Term Effects on Serum Parathyroid Hormone Concentration

W. Probst, J. A. Fischer and U. Binswanger

Section of Nephrology and Research Laboratory for Calcium Metabolism, Departments of Internal Medicine and Orthopedic Surgery, University of Zürich, Switzerland

Abstract. Eleven patients on chronic haemodialysis treatment thrice weekly received 1 µg 1,25(OH)₂D3 i.v. after each dialysis for 3 weeks. Phosphate binders were mainly CaCO₃, supplemented in a few patients by moderate amounts of Al(OH)₃. Ionised calcium was measured by ion-selective electrode, normal values being 1.28–1.42 mmol/l. PTH was estimated by an N-terminal-sensitive assay; normal values are <0.25 ng/ml. Results before and after 1,25(OH)₂D3 were: ionised calcium before haemodialysis, 1.19 ± 0.12 and 1.17 ± 0.14; ionised calcium after haemodialysis, 1.33 ± 0.07 and 1.30 ± 0.09; PTH before haemodialysis, 1.39 ± 0.71 and 1.38 ± 0.69; PTH after haemodialysis, 0.64 ± 0.22 and 0.60 ± 0.17; Phosphate before haemodialysis, 1.85 ± 0.48 and 2.18 ± 0.43 (*P* < 0.05). No change of PTH concentration and ionised calcium before and after haemodialysis treatment could be documented after i.v. 1,25(OH)₂D3 treatment. Mild and severe hyperparathyroidism were indistinguishable. Increased serum calcium concentrations therefore appear to be required for the suppression of PTH secretion by i.v. 1,25(OH)₂D3 therapy.

Key words: Haemodialysis; i.v. 1,25(OH)₂ vitamin D3; Serum parathyroid hormone

Introduction

Excessive parathyroid hormone (PTH) secretion is an important cause of renal osteodystrophy [1]. Serum PTH concentrations are increased early in the course of renal disease and are linked initially to predominantly intracellular phosphate retention [2]. This is followed at later stages of the disease by an extracellular increase of inorganic phosphate [3]. Phosphate retention is related to reduced ionised calcium concentrations, suppressed synthesis of 1,25(OH)₂ vitamin D3 (1,25(OH)₂D3) and the expression of its receptors. All these factors participate in the aetiology of renal hyperparathyroidism [4].

Suppression of parathyroid hormone secretion in renal disease is related to elevated serum ionised calcium concentrations [5]. Normal serum PTH values, however, are difficult to reach and are usually only observed in hypercalcaemic patients [6]. This was true with calcium supplementation as well as with the peroral administration of vitamin D metabolites [7]. A new approach for the treatment of renal hyperparathyroidism was initiated by the group of Slatopolsky with the use of intravenously administered 1,25(OH)₂D3 [8]. The authors postulated that a more pronounced suppression of parathyroid hormone secretion was brought about with intravenous 1,25(OH)₂D3 than with its peroral administration [9].

The present study was carried out to differentiate between effects of intravenous 1,25(OH)₂D3 and the increase in serum ionised calcium before and during haemodialysis treatment. Low doses of intravenously

administrated $1,25(\text{OH})_2\text{D}_3$ did not change serum calcium values at the start of haemodialysis; treatment-related augmentation of ionised calcium suppressed PTH to the same extent before and after i.v. $1,25(\text{OH})_2\text{D}_3$.

Patients and Methods

The study was carried out in 11 patients on chronic haemodialysis, mean age 36.2 ± 14.2 (SD) years, treated thrice weekly for 2.5–3.5 h by means of high-flux polysulphone filters (Fresenius F60), using acetate dialysate with 1.5 mmol/l of calcium (Table 1). One microgram of $1,25(\text{OH})_2\text{D}_3$ (Abbott Company) was administered i.v. after each dialysis session for 3 weeks. Serum ionised calcium by ion-selective electrode (AVL company, CV=0.7%), immunoreactive N-terminal parathyroid hormone (PTH) (CV=14%) [10] and inorganic phosphorus (ammonium molybdate reaction, CV=1.8%) were measured before and after i.v. $1,25(\text{OH})_2\text{D}_3$ treatment. Moreover, effects of individual haemodialysis treatments on the increase of the serum ionised calcium and the suppression of PTH were evaluated. All the patients were kept on calcium carbonate as a phosphate-binding agent, supplemented in some candidates with small doses of aluminium hydroxide. Mean serum aluminium values were increased 5-fold as compared to normal subjects, but distributed within the Gaussian distribution of the haemodialysis population. Statistical analysis was performed by Student's t-test for paired data.

Results (Fig. 1)

Before treatment with $1,25(\text{OH})_2\text{D}_3$, the serum ionised calcium values amounted to 1.19 ± 0.12 mmol/l (normal 1.28–1.42 mmol/l, mean ± 2 SD). During individual haemodialysis treatments, serum ionised calcium was increased to 1.33 ± 0.7 mmol/l. Corresponding values, after 3 weeks of i.v. $1,25(\text{OH})_2\text{D}_3$ treatment were 1.17 ± 0.14 mmol/l and 1.30 ± 0.09 mmol/l respectively. N-terminal PTH concentrations were 1.39 ± 0.71 ng/ml and 0.64 ± 0.22 ng/ml before and after dialysis prior to $1,25(\text{OH})_2\text{D}_3$ treatment (normal <0.25 ng/ml). After 3 weeks i.v. $1,25(\text{OH})_2\text{D}_3$ PTH amounted to 1.38 ± 0.69 ng/ml before and 0.60 ± 0.17 ng/ml after haemodialysis. Serum phosphorus concentrations amounted to 1.85 ± 0.48 mmol/l (normal 0.7–1.6 mmol/l) before $1,25(\text{OH})_2\text{D}_3$ treatment. After $1,25(\text{OH})_2\text{D}_3$ the values were slightly increased to 2.18 ± 0.43 mmol/l ($P < 0.05$). On correlation analysis the decrease of PTH was related

to the increase of serum ionised calcium during haemodialysis before as well as after $1,25(\text{OH})_2\text{D}_3$ treatment (Fig. 2).

Discussion



Increased serum ionised calcium concentrations are known to suppress PTH secretion [1]. In chronic renal insufficiency with and without dialysis treatment, near normal PTH values have only been observed during hypercalcaemia. This observation was related by Brown and others to an increased set point for PTH secretion in hyperplastic parathyroid glands [11–13]. Besides calcium supplementation, peroral vitamin D and its metabolites reduced serum PTH in concert with increased calcium values. Reduced serum PTH values were anticipated from in-vitro results, revealing suppression of mRNA values encoding for pre-pro-PTH by $1,25(\text{OH})_2\text{D}_3$ independent of ambient calcium concentrations [11,12]. Several other groups showed suppression of PTH by $1,25(\text{OH})_2\text{D}_3$ in vitro [14–16].

In contrast, direct suppression of the PTH secretion by $1,25(\text{OH})_2\text{D}_3$ is questionable in vivo and has not been universally demonstrated [7]. With peroral vitamin D metabolites a direct suppression of PTH secretion could not be dissociated from concomitantly increased serum calcium concentrations. An alternative arose when $1,25(\text{OH})_2\text{D}_3$ administered intravenously was claimed to suppress PTH secretion directly [8]. The mechanism was thought to be increased sensitivity to calcium-induced suppression of PTH secretion [9]. In these studies calcium was increased within minutes of i.v. $1,25(\text{OH})_2\text{D}_3$ administration and almost simultaneously, PTH decreased. The two variables, increased serum concentrations of $1,25(\text{OH})_2\text{D}_3$ and calcium, have not been clearly separated. None the less, increased serum calcium values in concert with or due to i.v. $1,25(\text{OH})_2\text{D}_3$ are effective in reducing serum PTH.

With the present study a potential direct suppression of PTH secretion by i.v. $1,25(\text{OH})_2\text{D}_3$ in the absence of increased ionised calcium concentration was investigated. Moreover, reduction of serum PTH levels as a consequence of haemodialysis treatment increasing serum calcium concentrations was confirmed [9]. Serum ionised calcium concentrations before the start of haemodialysis were stable and in a slightly hypocalcaemic range; $1 \mu\text{g}$ $1,25(\text{OH})_2\text{D}_3$ administered three times weekly by the i.v. route for 3 weeks did not alter mean serum PTH values in our patients. Moreover, grouping of the patients with great and small initial PTH values did not affect our conclusions. The quantity of phosphate binders administered was left unchanged. As a result of the treatment with $1,25(\text{OH})_2\text{D}_3$ serum inorganic phosphorus values were slightly elevated ($P < 0.05$). This observation reveals a

Table 1. Patients' characteristics and biochemical data before i.v. 1,25(OH)₂D₃ treatment

Patient	Age	Sex	Diagnosis	Duration of renal insufficiency (years)	Duration of dialysis therapy (months)	Ionised serum calcium (mmol/l)	Serum alkaline phosphatase (U/l)	Serum inorganic phosphate (mmol/l)	N-terminal PTH (ng/ml)
1	21	m	Rapidly progressive GN	01.0	13	1.25	49	2.07	0.33
2	33	f	Reflux nephropathy	05.0	36	0.91	76	2.02	2.42
3	30	f	Unknown origin	03.0	16	1.24	102	2.11	0.97
4	30	f	Lupus nephritis	02.5	16	1.11	45	2.02	1.27
5	24	m	Urethral valves	13.0	14	1.09	118	1.35	1.99
6	63	m	Analgesic nephropathy	10.0	02	1.20	89	1.69	2.33
7	21	m	Adult polycystic disease	08.0	39	1.20	100	0.99	1.11
8	53	m	Adult polycystic disease	01.0	14	1.21	54	1.59	2.00
9	48	f	Glomerulonephritis	16.0	23	1.35	105	1.53	0.79
10	28	f	Mesangial proliferative GN	06.0	11	1.25	61	2.39	1.50
11	47	m	Diabetic nephropathy	04.0	24	1.33	34	2.64	0.63
			Normal values			1.28–1.42	30–115	0.7–1.16	<0.25

Fig. 1. Serum ionised calcium, PTH and phosphorus before and after i.v. 1,25(OH)₂D₃. Effects of haemodialysis treatment.Fig. 2. Correlation between changes in ionised serum calcium and PTH during haemodialysis treatment before and after 1,25(OH)₂D₃.

biological effect of the 1,25(OH)₂D₃ administered and perhaps can be considered as a contribution to the lack of suppressive effect of i.v. administered 1,25(OH)₂D₃ on PTH, as recently suggested by Slatopolsky et al [17]. In contrast to this opinion Brandi et al [18] showed a suppression of PTH with i.v. 1- α -hydroxyvitamin D₃ despite a marked increase of serum phosphorus; the suppression of PTH, however, was accompanied by a marked increase in the serum calcium concentration.

The 1,25(OH)₂D₃ dose used was taken from published data which showed suppression of PTH in some patients [8]. This was accentuated during prolonged treatment with increasing doses of 1,25(OH)₂D₃ concomitant with an increase of the serum ionised calcium concentrations. More prolonged treatment with 1 μ g of 1,25(OH)₂D₃ as used in the present study might suppress PTH secretion in the absence of increased serum calcium concentrations. Serum ionised calcium was always elevated during haemodialysis treatment, and PTH reduced accordingly. Here also, the values were indistinguishable before and after 1,25(OH)₂D₃ treatment. With a larger dose of 1,25(OH)₂D₃ during a prolonged treatment time and while decreasing the dialysate calcium to 2.5 mEq/l,

Dunlay et al [19] showed an effect on PTH without a significant increase in serum calcium concentration. The sigmoidal shape relation between PTH and haemodialysis-induced hypo- and hypercalcaemia was displaced, showing a smaller PTH stimulation at low serum calcium concentrations after 1,25(OH)₂D₃; pre- and post-treatment relations, however, were only marginally affected. With a long-term intermittent treatment by i.v. calcitriol inducing an increase of the serum calcium concentration, an amelioration of osteitis fibrosa in dialysis patients was recently demonstrated by bone histomorphometry [20].

In conclusion, short-term i.v. administration of 1,25(OH)₂D₃ in slightly hypocalcaemic haemodialysis patients did not reduce serum PTH concentrations. Suppression of serum PTH by haemodialysis-induced increases of the ionised serum calcium concentrations was the same before and after i.v. 1,25(OH)₂D₃ treatment. These data might suggest a synergistic effect of 1,25(OH)₂D₃ and the increase in serum ionised calcium concentration for the suppression of serum PTH concentrations in haemodialysis patients.

Acknowledgements. This work was supported by the Swiss National Science Foundation grant 3.924-0.87 and the Canton of Zurich. Support by the staff of the haemodialysis unit, the laboratory technicians, and the secretary Mrs Cajet is appreciated.

References

1. Arnaud C. Hyperparathyroidism and renal failure. *Kidney Int* 1973; 4: 89-95
2. Binswanger U, Dambacher MA, Fischer JA. Sekundärer renaler Hyperparathyreoidismus. In: Hauri D, Schmucki O, ed. *Erkrankungen der Nebenschilddrüsen und Nebennieren*. Gustav Fischer Verlag, Stuttgart, 1985: 38-47
3. Lee DBN, Goodman WG, Coburn JW. Renal osteodystrophy. Some new questions on an old disorder. *Am J Kidney Dis* 1988; 5: 365-376
4. Feinfeld DA, Sherwood LM. Parathyroid hormone and 1,25(OH)₂D₃ in chronic renal failure. *Kidney Int* 1988; 33: 1049-1058
5. Meyrier A, Marsac J, Richet G. The influence of a high calcium carbonate intake on bone disease in patients undergoing haemodialysis. *Kidney Int* 1973; 4: 146-153
6. Slatopolsky E, Weerts C, Lopez-Hilker S et al. Calcium carbonate as a phosphate binder in patients with chronic renal failure undergoing dialysis. *N Engl J Med* 1986; 315: 157-161
7. Binswanger U, Fischer JA, Iselin H et al. 1,25-dihydroxycholecalciferol treatment of clinically asymptomatic renal osteodystrophy. *Miner Electrolyte Metab* 1979; 2: 103-115
8. Slatopolsky E, Weerts C, Thielan J, Horst R, Harter H, Martin KJ. Marked suppression of secondary hyperparathyroidism by intravenous administration of 1,25-dihydroxycholecalciferol in uremic patients. *J Clin Invest* 1984; 74: 2136-2143
9. Delmez JA, Tindira C, Grooms P, Dusso A, Windus DW, Slatopolsky E. Parathyroid hormone suppression by intravenous 1,25-dihydroxyvitamin D: A role for increased sensitivity to calcium. *J Clin Invest* 1989; 83: 1349-1355
10. Born W, Dambacher MA, Meyrier A, Ardaillou R, Fischer JA. Parathyroid suppressibility in hyperparathyroidism due to chronic renal failure: studies with autotransplanted parathyroid tissue. *Clin Endocrinol* 1982; 17: 333-343
11. Russel J, Lettieri D, Sherwood LM. Suppression by 1,25(OH)₂D₃ of transcription of the parathyroid hormone gene. *Endocrinology* 1986; 119: 2864-2866
12. Silver J, Naveh-Many T, Mayer H, Schmelzer HJ, Popovtzer MM. Regulation by vitamin D metabolites of parathyroid hormone gene transcription in vivo by the rat. *J Clin Invest* 1986; 78: 1296-1301
13. Brown EM, Wilkson RE, Eastman RC, Pallotta J, Marynick SP. Abnormal regulation of parathyroid hormone release by calcium in secondary hyperparathyroidism due to chronic renal failure. *J Clin Endocrinol Metab* 1982; 54: 172-179
14. Chan YL, McKay C, Dye E, Slatopolsky E. The effect of 1,25-dihydroxycholecalciferol on parathyroid hormone secretion by monolayer cultures of bovine parathyroid cells. *Calcif Tissue Int* 1986; 38: 27-32
15. Chertow BD, Baylink DJ, Wergedal JE, Su MH, Norman AW. Decrease in serum immunoreactive parathyroid hormone in rats and in parathyroid hormone secretion in vitro by 1,25-dihydroxycholecalciferol. *J Clin Invest* 1975; 56: 668-678
16. Au WYW, Bukowski A. Inhibition of parathyroid hormone secretion by vitamin D metabolites in organ cultures of rat parathyroids. *Fed Proc* 1976; 35: 530 (Abstr)
17. Slatopolsky E, Lopez-Hilker S, Dusso A, Morrissey JJ, Martin KJ. Parathyroid hormone secretion: Perturbations in chronic renal failure. In: De Broe ME, Van de Vyver FL, ed. *Bone and Renal Failure*, Contr Nephrol, Karger Basel 1988; 64: 16-24
18. Brandi L, Dagaard H, Tvedegaard E, Storm T, Olgaard K. Effect of intravenous 1- α -hydroxyvitamin D₃ on secondary hyperparathyroidism in chronic uremic patients on maintenance hemodialysis. *Nephron* 1989; 53: 194-200
19. Dunlay R, Rodriguez M, Felsenfeld AJ, Llach F. Direct inhibitory effect of calcitriol on parathyroid function (sigmoidal curve) in dialysis. *Kidney Int* 1989; 36: 1093-1098
20. Andress DL, Norris KC, Coburn JW, Slatopolsky EA, Sherrard DJ. Intravenous calcitriol in the treatment of refractory osteitis fibrosa of chronic renal failure. *N Engl J Med* 1989; 321: 274-279

Received for publication 12.10.89

Accepted in revised form 26.2.90